

Abstract No. zhou673

**High resolution structure of a potassium channel – antibody Fab complex**

Y. Zhou, J.H. Morais-Cabral, A. Kaufman, R. Mackinnon (HHMI, Rockefeller University)

Beamline: X25

**Introduction:** Potassium channels control the electric potential across cell membranes by catalyzing the rapid, selective diffusion of  $K^+$  ions down their electrochemical gradient<sup>1</sup>. The process of dehydration, transfer and re-hydration of a  $K^+$  ion is catalyzed by the channel's selectivity filter. To study the mechanism of ion translocation in potassium channels, it is necessary to solve the  $K^+$  channel structure at a high resolution that would reveal protein chemistry and the ordered water molecules around  $K^+$  ions with high accuracy.

**Methods and Materials:** We raised monoclonal antibody against the KcsA potassium channel, and purified the Fab fragment. A KcsA-Fab complex with a stoichiometry of one Fab fragment per channel subunit was produced and crystallized in space group I4. Data collections were conducted at X25 on two types of frozen crystals: KcsA-Fab complex crystallized in high concentration of  $K^+$  ions diffracted X-rays to 2.0 Å (cell:  $a = 155.33$ ,  $b = 155.33$ ,  $c = 76.27$ ,  $\alpha = \beta = \gamma = 90^\circ$ ), and the complex crystallized in low concentration of  $K^+$  ions diffracted to 2.3 Å (cell:  $a = 155.29$ ,  $b = 155.29$ ,  $c = 75.74$ ,  $\alpha = \beta = \gamma = 90^\circ$ ). Phases were solved by molecular replacement<sup>2,3</sup>. The high- $K^+$  structure was refined to  $R_f$  and  $R_w$  of 23.3% and 21.8%, and the low- $K^+$  structure was refined to  $R_f$  and  $R_w$  of 23.5% and 21.8%, respectively.

**Results:** In these two structures, ions within the selectivity filter were well resolved. The structures also revealed ordered water molecules around the  $K^+$  ions near the entryways to the filter. These two structures allowed us to address the mechanisms of  $K^+$  ion hydration and dehydration, and the response of the selectivity filter the changing ionic environment imposed by channel gating.

**Acknowledgments:** We thank the staff at the NSLS X25 and CHESS A1 and F1 for assistance. This study was supported by a grant from the National Institutes of Health to R.M. R.M. is an investigator in the Howard Hughes Medical Institute.

**References:**

- [1] B. Hille, "Ionic Channels of Excitable Membranes" (Sinauer, Sunderland, 1992).
- [2] D.A. Doyle *et al.* "The structure of the potassium channel: molecular basis of  $K^+$  conduction and selectivity". *Science* 280, 69-77 (1998).
- [3] B. Braden *et al.* "Three-dimensional structures of the free and the antigen-complexed Fab from monoclonal anti-lysozyme antibody D44.1". *J. Mol. Biol.* 243, 767-781 (1994).